

# DESERT NEWS.

SALT LAKE CITY, UTAH TERRITORY, TUESDAY EVENING, AUGUST 30, 1881. NO. 235.

**THE DESERT NEWS.**  
Published every evening except Sunday.  
One copy, one year, with Postage, \$10.00  
" " six months, " 5.00  
" " three months, " 2.50  
**DESERET NEWS.**  
SEMI-WEEKLY.  
Published every Tuesday.  
One copy, one year, with Postage, \$5.00  
" " six months, " 2.50  
" " three months, " 1.25  
**DESERET NEWS.**  
WEEKLY.  
Published every Wednesday.  
One copy, one year, with Postage, \$3.00  
" " six months, " 1.50  
" " three months, " .75  
**TERMS - IN ADVANCE.**  
OFFICE - Corner of South and East Temple Streets.

The News Book and Job Printing Office  
Has superior facilities for the execution of  
All kinds of  
Book and Job Printing.

BOOK-BINDING & PAPER RULING

**NEW YORK TRADE.**

**HEMPHILL, HAMLIN & Co.**  
CARPETING,  
FLOOR OILS, BRUSSETS, MATS, ETC.  
344 N. 3rd Broadway,  
NEW YORK.

Our Goods are kept by Z. C. M. I. and its Branches.

**THOS. M. ARGALL & Co.**  
Manufacturers of  
**MEN'S & BOYS'**  
**CLOTHING.**  
330 Broadway,  
NEW YORK.

BRINCKHOFF, TURNER & Co.,  
100 Duane Street, New York.

Manufacturers of and Dealers in  
COTTON, WOOL, LINEN, and other fabrics,  
Hosiery, Trimmings, and other notions,  
Ducks for overalls, blue, brown, or in the grey,  
COTTON CANVAS of all numbers,  
14 to 16 inches wide for Decks, Caps, Trunks,  
and Wagon Coverings. Machine Aprons and  
other purposes, constantly in store and made to order.  
For U. S. Shipping Co. "Standard"  
and "Eagle" by the case or 500 garments.  
For these goods can be ordered at Z. C. M. I.

**B. W. MERRIAM & Co.**  
MANUFACTURERS OF AND DEALERS IN  
**LOOKING GLASSES**  
OF EVERY DESCRIPTION.  
NO. 577 Broadway,  
NEW YORK.

For sale by Z. C. M. I.

**BENEDICT, HALL & Co.**  
Wholesale Manufacturers of  
**BOOTS**  
-AND-  
**SHOES.**  
Nos. 384 and 386 Grand Street,  
New York.

**BATES & JOHNSON.**  
SUCCESSORS TO  
WYLLIS H. WARNER,  
HIGH AND LOW PASSBROS.

**STEAM HEATING APPARATUS.**  
114 Leonard Street, N. Y.

David James, Agent, Salt Lake City.

See the Assembly Hall, Salt Lake City,  
is warmed by our Apparatus erected by Mr. James.

**PECK BROS. & Co.**  
MANUFACTURERS OF  
BRASS AND PLATED WORK FOR WATER GAS & STEAM  
JOURNALS IN  
**PLUMBERS MATERIALS.**  
75 BREKMAN STREET,  
NEW YORK.

Order by permission to David James, Salt Lake City.

**A. W. FABER'S**  
STATIONERY ARTICLES & ARTISTS' MATERIALS.  
MAY BE HAD AT  
**Z. C. M. I.**  
Gold Pens and Pencil Cases, Pen-  
holders, Rubber Goods, Lead Pencil  
Cases, Etc.

**USE**  
**CLARK'S**  
**TRADE**  
**MARK**  
**ON**  
**SPPOOL COTTON.**

**GEORGE A. CLARK,**  
SOLE AGENT.

The BEST and MOST POPULAR  
Sewing Thread of Modern Times.

Beware of IMITATIONS.

A COMPLETE ASSORTMENT  
FOR SALE BY  
**Z. C. M. I.**  
Branches and Dealers Everywhere.

**KEITH BROTHERS,**  
MANUFACTURERS AND JOINERS OF  
**Hats, Caps,**  
**FURS**  
**AND**  
**STRAW GOODS.**  
**BUCK GOODS, ETC.**  
246, 248, 250 & 252, Madison Street,  
CHICAGO, ILL.  
AN ARTICLE OF WORLD-WIDE RENOWN  
IMPORTERS PURE REFINED LARD.

TRY IT!  
**N. K. FAIRBANK & Co.**  
CHICAGO.

For Sale by Z. C. M. I. and its Branches.

**JOHN C. NEEMES & Co.**  
MANUFACTURERS

Our Goods are kept by Z. C. M. I. and its Branches.

**CONFECTIONERS**  
Y. 34, 36, 38 & 40 MICHIGAN ST.

CHICAGO.

Our Goods are kept by Z. C. M. I. and its Branches.

**TURNER & RAY,**  
DEALERS IN

**LEATHER AND FINDINGS**

**ROOT AND SHOES MANUFACTURERS**  
TURNER'S GOODS, MANHATTAN,  
N. Y. TOOLS, ETC., ETC.

Wholesale Manufacturers Goods and Machinery a Specialty.  
CHICAGO, ILL.

**ST. LOUIS TRADE.**

**Leggett & Myers Tobacco Co.**  
MANUFACTURERS OF THE  
JAB-BANDER, COMMERCIAL, CLIPPED, AMERICAN,  
AND OTHER BRANDS OF  
**TOBACCO.**  
In sale by Z. C. M. I. and Branches Everywhere.

**L. M. RUMSEY & Co.**  
Manufacturers and Jobbers of  
**PUMPS AND FIRE ENGINES,**  
and Pipe and Sheet Lead, Cast Pipe,  
Flanges and Steam Fitters.  
Brass Goods, Belling,  
and other work.  
10 NORTH SECOND ST., ST. LOUIS, MO.

**AGRICULTURAL IMPLEMENTS,**  
and Wire, Barbed Wire, Fencing,  
Etc., Etc.

**Fairbanks' Scales!**  
THE STANDARD OF THE WORLD.

**Eclipse Wind Mills.**  
The Strongest Mills Made.

**SAFEST TO BUY.**  
Because warranted the best (same as Scales)  
and because it is substantial. Contains all  
improvements. Prices lowest, quality superior.  
See the catalogue describing article within.

**FAIRBANKS & Co.,**  
102 & 104 WASHINGTON ST., ST. LOUIS, MO.

AGENTS FOR  
**The Hancock Inspirator.**  
THE BEST BOILER FEEDER KNOWN.

**Southern Company**  
STRICTLY PURE WHITE LEAD.

THE SOUTHERN WHITE LEAD

STRICTLY PURE.

For sale by Z. C. M. I. and its Branches.

**THE SOUTHERN WHITE LEAD**

STRICTLY PURE.

For sale by Z. C. M. I. and its Branches.

**THE SOUTHERN WHITE LEAD**

STRICTLY PURE.

For sale by Z. C. M. I. and its Branches.

**THE SOUTHERN WHITE LEAD**

STRICTLY PURE.

For sale by Z. C. M. I. and its Branches.

**THE SOUTHERN WHITE LEAD**

STRICTLY PURE.

**BY TELEGRAPH.**

**AMERICAN.**

**THE PRESIDENT'S CONDITION.**

**The Name.**

**EXECUTIVE MANSION, 8.15 p.m.**

The daily rise of the President's temperature began later this afternoon than yesterday, and rose eight-tenths of a degree higher. The frequency of his pulse is not the same as at this hour yesterday. It has been falling during the day, and besides, during the morning, the pulse was 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

The pulse was 110, temperature 100.6, and during the evening, 100.6.

porter made use of any expressions beyond the copy left he should not record him a second interview. The reporter then asked, "Are you re-portedly correct in an evening paper to-day, that you had no doubt of his recovery?"

Dr. Hamilton: "No, sir, I would not say that of any patient, and much less of the President, in whose case it is particularly difficult to make a prognosis. The reporter misinterpreted me, and I said I would not say that, but that I believe it is not absolute assurance. It is simply my faith."

**Blame to Lowell.**

Washington, 29.—At half-past 10 to-night the general opinion of the President's condition was that the afternoon pulse rose to 112 and temperature to 108, which was a little higher than the surgeon's expected. The pulse has now fallen to 108, fever subsiding, and the parotid swelling is steadily improving or, at least, diminishing in size. Apprehensions of serious blood poisoning grow less every hour.

**Not Unfavorable.**

Executive Mansion, 8.15 p.m.—The evening bulletin is generally accepted here as a confirmation of the hopes of the day. The President's fever, which was a little higher to-night than last night, has begun to subside, and all symptoms continue satisfactory. Dr. Agnew, as he was leaving the Mansion, was asked what he thought of the bulletin. "It is a very good one," he said, "and a fair and accurate statement of the President's condition." "And what view do you take of his condition?" "There is nothing unfavorable about it at all."

**Consolidated.**

10 p.m.—The improvement in the President's condition, which began Saturday, has continued without interruption to the present. The attending surgeons agree that the case shows a marked and decided improvement. The President's condition, which holds out promise of permanency. The President's attendants take even a more sanguine view of the case, and hope for a recovery within a few days, which verges upon assured confidence. Even the President himself feels this strong current of renewed hope, and his reinvigorated mental energy co-operates with the physical. As stated in the early morning bulletin, the patient had better night than at any time before. The middle of last week, when he was still broken at rather short intervals, but was more natural and peaceful than heretofore, with less disturbance from the parotid gland. He was sponged off with alcohol and water every three or four hours, and swallowed liquid nourishment at intervals of two or three hours. The night, the morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back, however, to 104. The patient rested quietly throughout the morning and night, and was able to eat a little. The morning examination, and the evening examination, all showed a decided improvement. Another slight incision made at the morning dressing in the right cheek gave the patient some relief from the pain, and the swelling had previously been 100 to 120. It soon fell back,